

# MAXIMUM AND MINIMUM VALUES

Math 130 - Essentials of Calculus

6 November 2019

## DEFINITIONS - REVIEW

## DEFINITIONS - REVIEW

## DEFINITION

Let  $c$  be a number in the domain of a function  $f$ . Then  $f(c)$  is the

- *absolute maximum value of  $f$  on the domain if  $f(c) \geq f(x)$  for all  $x$  in the domain.*

## DEFINITIONS - REVIEW

## DEFINITION

Let  $c$  be a number in the domain of a function  $f$ . Then  $f(c)$  is the

- absolute maximum value of  $f$  on the domain if  $f(c) \geq f(x)$  for all  $x$  in the domain.
- absolute minimum value of  $f$  on the domain if  $f(c) \leq f(x)$  for all  $x$  in the domain.

## DEFINITIONS - REVIEW

## DEFINITION

Let  $c$  be a number in the domain of a function  $f$ . Then  $f(c)$  is the

- absolute maximum value of  $f$  on the domain if  $f(c) \geq f(x)$  for all  $x$  in the domain.
- absolute minimum value of  $f$  on the domain if  $f(c) \leq f(x)$  for all  $x$  in the domain.

These values are also called the extreme values of  $f$ .

## DEFINITIONS - REVIEW

### DEFINITION

Let  $c$  be a number in the domain of a function  $f$ . Then  $f(c)$  is the

- absolute maximum value of  $f$  on the domain if  $f(c) \geq f(x)$  for all  $x$  in the domain.
- absolute minimum value of  $f$  on the domain if  $f(c) \leq f(x)$  for all  $x$  in the domain.

These values are also called the extreme values of  $f$ .

A function often has values that behave like maximum and minimum values if you only look at nearby input values. These are called local extrema.

### DEFINITION

The number  $f(c)$  is a

- local maximum value of  $f$  if  $f(c) \geq f(x)$  when  $x$  is near  $c$ .

## DEFINITIONS - REVIEW

### DEFINITION

Let  $c$  be a number in the domain of a function  $f$ . Then  $f(c)$  is the

- absolute maximum value of  $f$  on the domain if  $f(c) \geq f(x)$  for all  $x$  in the domain.
- absolute minimum value of  $f$  on the domain if  $f(c) \leq f(x)$  for all  $x$  in the domain.

These values are also called the extreme values of  $f$ .

A function often has values that behave like maximum and minimum values if you only look at nearby input values. These are called local extrema.


### DEFINITION

The number  $f(c)$  is a

- local maximum value of  $f$  if  $f(c) \geq f(x)$  when  $x$  is near  $c$ .
- local minimum value of  $f$  if  $f(c) \leq f(x)$  when  $x$  is near  $c$ .

## EXTREMA OF A FUNCTION

Consider the function  $f(x) = 3x^4 - 16x^3 + 18x^2$  on the domain  $-1 \leq x \leq 4$ . Where are the absolute maximum and absolute minimum values, and what are they? Are there any local minimum and local maximum values?





# EXTREMA OF A FUNCTION

Whether or not a function has absolute/local extrema depends on the function AND the domain you look at it on.

## EXTREMA OF A FUNCTION

Whether or not a function has absolute/local extrema depends on the function AND the domain you look at it on.

### EXAMPLE

- 1 *Considered on the domain  $(-\infty, \infty)$ ,  $f(x) = x^2$  has an absolute minimum, but nothing else.*

## EXTREMA OF A FUNCTION

Whether or not a function has absolute/local extrema depends on the function AND the domain you look at it on.

### EXAMPLE

- 1 *Considered on the domain  $(-\infty, \infty)$ ,  $f(x) = x^2$  has an absolute minimum, but nothing else.*
- 2 *Considered on the domain  $(-\infty, \infty)$ ,  $f(x) = x^3$  has no extrema at all!*

## EXTREMA OF A FUNCTION

Whether or not a function has absolute/local extrema depends on the function AND the domain you look at it on.

### EXAMPLE

- 1 *Considered on the domain  $(-\infty, \infty)$ ,  $f(x) = x^2$  has an absolute minimum, but nothing else.*
- 2 *Considered on the domain  $(-\infty, \infty)$ ,  $f(x) = x^3$  has no extrema at all!*

It turns out that if you consider a continuous function on a closed interval, of the form  $[a, b]$ , you're guaranteed an absolute maximum and minimum.

### THEOREM (THE EXTREME VALUE THEOREM)

*If  $f$  is continuous on a closed interval, then it always attains an absolute maximum value and an absolute minimum value on that interval.*

# LOCATING EXTREME VALUES

Observing some of the pictures we've had so far, the following theorem is apparent:

## THEOREM (FERMAT'S THEOREM)

*If  $f$  has a local maximum or minimum at  $c$ , and if  $f'(c)$  exists, then  $f'(c) = 0$ .*

## LOCATING EXTREME VALUES

Observing some of the pictures we've had so far, the following theorem is apparent:

### THEOREM (FERMAT'S THEOREM)

*If  $f$  has a local maximum or minimum at  $c$ , and if  $f'(c)$  exists, then  $f'(c) = 0$ .*

It's possible that a function could have a local extrema at a place where  $f'(c) \neq 0$ , for example, consider  $f(x) = |x|$ .

## LOCATING EXTREME VALUES

Observing some of the pictures we've had so far, the following theorem is apparent:

### THEOREM (FERMAT'S THEOREM)

*If  $f$  has a local maximum or minimum at  $c$ , and if  $f'(c)$  exists, then  $f'(c) = 0$ .*

It's possible that a function could have a local extrema at a place where  $f'(c) \neq 0$ , for example, consider  $f(x) = |x|$ . It turns out that what we're really looking for are *critical numbers*.

### DEFINITION (CRITICAL NUMBER)

*A critical number of a function  $f$  is a number  $c$  in the domain of  $f$  such that either  $f'(c) = 0$  or  $f'(c)$  does not exist.*

## CRITICAL NUMBERS - EXAMPLES

## EXAMPLE

Find the critical numbers of the following functions

①  $A(t) = t^3 - 6t^2 - 36t + 7$



## CRITICAL NUMBERS - EXAMPLES

## EXAMPLE

Find the critical numbers of the following functions

①  $A(t) = t^3 - 6t^2 - 36t + 7$

②  $f(x) = \sqrt[3]{x}$

## CRITICAL NUMBERS - EXAMPLES

## EXAMPLE

Find the critical numbers of the following functions

①  $A(t) = t^3 - 6t^2 - 36t + 7$

②  $f(x) = \sqrt[3]{x}$

③  $g(x) = x^3 + 3x^2 - 24$

# THE CLOSED INTERVAL METHOD

## THEOREM

*To find the absolute maximum and minimum values of a continuous function  $f$  on a closed interval  $[a, b]$ :*

- 1 Find the critical numbers of  $f$  in the interval  $(a, b)$  and compute the values of  $f$  at these numbers.*
- 2 Find the values of  $f$  at the endpoints of the interval.*
- 3 The largest of the output values from steps 1 and 2 is the absolute maximum value; the smallest of these values is the absolute minimum value.*

## FINDING EXTREMA

## EXAMPLE

Find the absolute maximum and absolute minimum values of  $f$  on the given interval.

①  $f(x) = x^3 - 3x + 1, [0, 3]$

②  $f(x) = x^3 - 6x^2 + 9x + 2, [-1, 4]$